

RESEARCH ARTICLE

Application of New Iterative Method and Adomian Decomposition Method to Hamel's Flow Problem

*Javed Ali¹¹Department of Mathematics & Statistics, Caledonian College of Engineering, Oman.

Received- 4 January 2017, Revised- 27 July 2017, Accepted- 24 October 2017, Published- 3 November 2017

ABSTRACT

In this work, we solve the Hamel's Problem using New Iterative Method (NIM) and Adomian Decomposition Method (ADM). This is a third order nonlinear boundary value problem. This problem arises when an incompressible fluid flows outward in a trough between two nonparallel plane walls. Application of these methods to this problem gives excellent numerical results with only three iterations. Moreover, accompanying residuals with these methods is an additional plus point which make these methods self-sufficient.

Keywords: Hamel's problem, Nonlinear boundary value problem, NIM, ADM, Incompressible fluid method.

1. INTRODUCTION

Higher order initial and boundary value problems have been investigated by many authors due to their physical importance and the potential for applications in hydrodynamic and hydro magnetic stability [1, 2]. Exact solutions for higher order differential equations are rare in literature. Researchers developed many methods to address this issue and to obtain approximate solutions of high accuracy. The most common techniques for obtaining a series form of the approximate solution, are; Adomian Decomposition Method (ADM) [3-5], variational iteration [6-8], homotopy perturbation [9-11], homotopy analysis [12], Differential Transform Method (DTM) [13] and optimal homotopy asymptotic method [14, 15].

A New Iterative Method (NIM) was introduced by Daftardar-Gejji and Jafari. It is an efficient technique to solve nonlinear functional equations [16].

Non-perturbation methods like ADM, NIM, DTM etc have certain advantages over routine numerical methods. Numerical methods use discretization which gives rise to rounding off errors causing loss of accuracy, and require large computer memory and time. On the other hand, the above mentioned methods yield analytical solutions and have certain advantages over standard numerical methods. These methods do not involve discretization of the variables and hence are free from rounding off errors and do not require large computer memory or time. In certain cases the obtained series solutions lead to the closed form solutions.

2. APPLYING NIM AND ADM

In this section we are not presenting general procedures of NIM and ADM. These methods have been extensively used in literature. We targeted these methods for the following boundary value problem [17]:

$$\frac{d^3y}{dx^3} + (2y+4)\frac{dy}{dx} = 0,$$

$$y(0) = 0, \quad y'(0.087) = 0, \quad y''(0) = -0.10.$$

$$\text{Let } L = \frac{d^3}{dx^3}, \quad \text{then } y = -L^{-1} \left\{ (2y+4)\frac{dy}{dx} \right\}, \quad \text{where } L^{-1} = -\int_0^x \int_0^x \int_0^x N dx \text{ and, } N = (2y+4)\frac{dy}{dx}.$$

To solve this problem, we set an algo according to the NIM and ADM. We seek the solution in the form

*Corresponding author. Tel.: +96899852897

Email address: javed.ali@caledonian.edu.om (J.Ali)

Double blind peer review under responsibility of DJ Publications

<http://dx.doi.org/10.18831/djcivil.org/2018011003>

2455-3581 © 2017 DJ Publications by Dedicated Juncture Researcher's Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

$$y = y_0 + \sum_{k=1}^{\infty} y_k \tag{1}$$

where y_0 is the solution of $Ly_0 = 0$, $y_0(0) = 0$, $y_0'(0.087) = 0$, $y_0''(0) = -0.10$, and y_i is obtained from

$$y_i = - \int_0^x \int_0^x \int_0^x N_{i-1} dx$$

At this stage, we decompose $N = \sum N_j$.

Now as,

$$N = \left(2 \sum_{i=0}^{\infty} y_i + 4 \right) \sum_{i=0}^{\infty} y_i = (2(y_0 + y_1 + y_2 + \dots) + 4)(y_0' + y_1' + y_2' + \dots),$$

so, in accordance to NIM & ADM, the following decomposition takes place as shown in table 1.

Table 1. Decomposition of NIM and ADM

NIM	ADM
$N_0 = 2y_0y_0' + 4y_0'$,	$N_0 = 2y_0y_0' + 4y_0'$,
$N_1 = 2y_0'y_1 + 2y_1'y_0 + 2y_1'y_1 + 4y_1'$,	$N_1 = 2y_0'y_1 + 2y_1'y_0 + 4y_1'$,
$N_2 = 2y_0'y_2 + 2y_1'y_2 + 2y_2'y_0 + 2y_2'y_1 + 2y_2'y_2 + 4y_2'$	$N_2 = 2y_0'y_2 + 2y_2'y_0 + 2y_1'y_1 + 4y_2'$
...	...

The components of the solution (1) are now given by

$$y_1 = - \int_0^x \int_0^x \int_0^x N_0 dt, \quad y_2 = - \int_0^x \int_0^x \int_0^x N_1 dt, \quad y_3 = - \int_0^x \int_0^x \int_0^x N_2 dt, \dots$$

Truncating the series solution (1), approximate solution is given by

$$\tilde{y} = y_0 + \sum_{k=1}^n y_k \tag{2}$$

Validity of the solution (2) can be confirmed by calculating residual at different mesh points of the domain. For NIM solution, the residual is

$$R_{NIM} = \frac{d^3}{dx^3}(\tilde{y}_{NIM}) + (2\tilde{y}_{NIM} + 4) \frac{d}{dx}(\tilde{y}_{NIM})$$

For ADM solution, the residual is

$$R_{ADM} = \frac{d^3}{dx^3}(\tilde{y}_{ADM}) + (2\tilde{y}_{ADM} + 4) \frac{d}{dx}(\tilde{y}_{ADM})$$

These residuals do not give the true error but they do indicate the validity of solution.

Now proceeding towards the solution we obtain, $\tilde{y} = y_0 + y_1 + y_2 + y_3 + O(x^{14})$.

$$\begin{aligned} \tilde{y}_{NIM} = & 0.0087x - 0.05x^2 - 0.0058x^3 + 0.0166604x^4 + 0.0012035x^5 - 0.00230135x^6 \\ & - 0.000135331x^7 + 0.000193158x^8 + 6.2927 \times 10^{-6}x^9 - 7.0407 \times 10^{-6}x^{10} \\ & - 7.3593 \times 10^{-7}x^{11} + 6.9008 \times 10^{-7}x^{12} + 4.6116 \times 10^{-8}x^{13} + O(x^{14}) \end{aligned}$$

$$\begin{aligned} \tilde{y}_{ADM} = & 0.0087x - 0.05x^2 - 0.0058x^3 + 0.0166604x^4 + 0.0012035x^5 - 0.00230135x^6 \\ & - 0.000135331x^7 + 0.000193158x^8 + 6.2927 \times 10^{-6}x^9 - 7.2169 \times 10^{-6}x^{10} \end{aligned}$$

$$-5.3134 \times 10^{-8}x^{11} + 5.0854 \times 10^{-8}x^{12} + 1.0844 \times 10^{-10}x^{13} + O(x^{14})$$

The comparison of numerical results is shown in table 2.

Table 2.Comparison of the numerical results

x	\tilde{y}_{NIM}	\tilde{y}_{ADM}	R_{NIM}	R_{ADM}
0.00	0.000	0.000	0.000	0.000
0.01	0.000081994	0.000081994	-3.0496×10^{-15}	-3.0496×10^{-15}
0.02	0.000153956	0.000153956	-1.9151×10^{-13}	-1.9167×10^{-13}
0.03	0.000215857	0.000215857	-2.1446×10^{-12}	-2.1469×10^{-12}
0.04	0.000267672	0.000267672	-1.1842×10^{-11}	-1.1859×10^{-11}
0.05	0.000309379	0.000309379	-4.4384×10^{-11}	-4.4459×10^{-11}
0.06	0.000340964	0.000340964	-1.3017×10^{-10}	-1.3042×10^{-10}
0.07	0.000362412	0.000362412	-3.2226×10^{-10}	-3.2296×10^{-10}
0.08	0.000373716	0.000373716	-7.0476×10^{-10}	-7.0440×10^{-10}

In figure 1, residual of the NIM is represented by the red solid curve and residual of the ADM is represented by the dotted curve.




Figure 1.Red solid curve - NIM residual & dotted curve - ADM residual

3. CONCLUSION

We used NIM and ADM for the solution of Hamel’s problem. Both the methods are easily applicable to higher order initial or boundary value problems and produce excellent results without any assumption and restrictions. Validity of the solutions was confirmed by calculating residuals at the mesh points. Both the methods are in complete agreement with each other.

REFERENCES

- [1] S.Chandrasekhar, Hydrodynamic Hydromagnetic Stability, Oxford University Press, London, 1961, pp. 1-641.
- [2] R.P.Agarwal, Boundary Value Problems for Higher Order Differential Equations, World Scientific, Singapore, 1986.
- [3] G.Adomian, Solving Frontier Problems of Physics: The Decomposition Method, Springer Science & Business Media, Germany, 2013.
- [4] G.Adomian, A Review of the Decomposition Method and some recent Results for Nonlinear Equations, Computers & Mathematics with Applications, Vol. 21, No. 5, 1991, pp. 101-127, [https://doi.org/10.1016/0898-1221\(91\)90220-X](https://doi.org/10.1016/0898-1221(91)90220-X).

- [5] A.M.Wazwaz, Approximate Solutions to Boundary Value Problems of Higher Order by the Modified Decomposition Method, *Computers & Mathematics with Applications*, Vol. 40, No. 6-7, 2000, pp. 679-691, [https://doi.org/10.1016/S0898-1221\(00\)00187-5](https://doi.org/10.1016/S0898-1221(00)00187-5).
- [6] Ji-Huan He, Variational Approach to the Sixth-Order Boundary Value Problems, *Applied Mathematics and Computation*, Vol. 143, No. 2, 2003, pp. 537-538, [https://doi.org/10.1016/S0096-3003\(02\)00381-8](https://doi.org/10.1016/S0096-3003(02)00381-8).
- [7] Shahid S.Siddiqi, Ghazala Akram and Imran Zulfiqar, Solution of Eleventh Order Boundary Value Problems using Variational Iteration Technique, *European Journal of Scientific Research*, Vol. 30, No. 4, 2009, pp. 505-525.
- [8] Muhammad Raheel Mohyuddin, Samra and Syed Mohammad Rizwan, Perturbation Unsteady Flows of 1-D Fluid, *Journal of Advances in Civil Engineering*, Vol. 1, No. 1, 2015, pp. 7-10, <http://dx.doi.org/10.18831/djccivil.org/2015011002>.
- [9] Ji-Huan He, Homotopy perturbation technique, *Computer Methods in Applied Mechanics and Engineering*, Vol. 178, No. 3, 1999, pp. 257-262, [https://doi.org/10.1016/S0045-7825\(99\)00018-3](https://doi.org/10.1016/S0045-7825(99)00018-3).
- [10] Ji-Huan He, Homotopy Perturbation Method for Solving Boundary Value Problems, *Physics Letters A*, Vol. 350, 2006, pp. 87-88, <https://doi.org/10.1016/j.physleta.2005.10.005>.
- [11] Jamshaid ul Rahman, Muhammad Raheel Mohyuddin, S.V.Satyanarayana, Syed Zahoor and Amira Al-Ghareebhi, Mathematical and Simulink Model for Paint Industry Effluent, *DJ Journal of Engineering Chemistry and Fuel*, Vol. 1, No. 2, 2016, pp. 1-8, <http://dx.doi.org/10.18831/djchem.org/2016021001>.
- [12] Shijun Liao, *Beyond Perturbation: Introduction to the Homotopy Analysis Method*, CRC Press, Florida, 2004.
- [13] Siraj-Ul-Islam, Siraj-Ul-Haq and Javed Ali, Numerical Solution of Special 12th - Order Boundary Value Problems using Differential Transform Method, *Communications in Nonlinear Science and Numerical Simulation*, Vol. 14, No. 4, 2009, pp. 1132-1138, <https://doi.org/10.1016/j.cnsns.2008.02.012>.
- [14] Javed Ali, S.Islam, Sirajul Islam and Gulzaman, The Solution of Multipoint Boundary Value Problems by the Optimal Homotopy Asymptotic Method, *Computers & Mathematics with Applications*, Vol. 59, No. 6, 2010, pp. 2000-2006, <https://doi.org/10.1016/j.camwa.2009.12.002>.
- [15] Javed Ali, Saeed Islam, Hamid Khan and Gul Zaman, The Solution of A Parameterized Sixth Order Boundary Value Problem by the Optimal Homotopy Asymptotic Method, *Proceedings of the Romanian Academy, Series A*, Vol. 12, No. 3, 2011, pp. 167-172.
- [16] V.Daftardar-Gejji and H.Jafari, An Iterative Method for solving Nonlinear Functional Equations, *Journal of Mathematical Analysis and Applications*, Vol. 316, No. 2, 2006, pp. 753-763, <https://doi.org/10.1016/j.jmaa.2005.05.009>.
- [17] T.Y.Na, *Computational Methods in Engineering Boundary Value Problems*, Academic Press, New York, 1980.